

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:JGRO-3-138

[European Legal view on Termination of Pregnancy](#)

Published On: April 28, 2017 | Pages: 051 - 055

Author(s): Siniša Franji

Pregnancy, another state or gravidity, is the state of a woman who carries a new being. The fertilization and ends with childbirth. The fertilized egg develops and goes through several different stages of development: embryo and fetus. ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/jgro.000038](https://doi.org/10.17352/jgro.000038)

[Open Access](#) [Research Article](#) PTZAID:JGRO-3-137

[Evaluation of Subendometrial and Intramyometrial Blood Flow after Intravenous Tranexamic Acid for Prevention of Postpartum Hemorrhage in Vaginal Delivery: A Randomized Controlled Study](#)

Published On: April 25, 2017 | Pages: 046 - 050

Author(s): Alaa M Ismail, Ahmed M Abbas*, Mohamed A Shahat and Mohammed K Ali

Objective: The study aims to test the effect of tranexamic acid (TA) on uterine vasculature and blood loss after vaginal delivery. ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/jgro.000037](https://doi.org/10.17352/jgro.000037)

[Open Access](#) [Research Article](#) PTZAID:JGRO-3-136

[Role of Soluble FMS-Like Tyrosine Kinase \(SFLT-1\) /Placental Growth Factor \(Plgf\) Ratio as Prognostic Marker for Cases of Preeclampsia](#)

Published On: April 15, 2017 | Pages: 037 - 045

Author(s): Gasser M Elbishry, Ihab F Serag Eldin, Ahmed A ElShahawy*, Gihan E Hawwary and Ahmed M Riad,

Objective: The aim of this study is to identify the role of sFlt-1/PlGF ratio as a prognostic marker for cases of

preeclampsia. Patients and methods: The current study is a case control study that was conducted over 90 cases of primigravida patients, 24-34 weeks of gestation, randomly selected patients from outpatient clinic and ER of Ain Shams Maternity Hospital ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/jgro.000036

[Open Access](#) | [Research Article](#) | PTZAID:JGRO-3-133

[Human Papilloma Virus vaccine – awareness and acceptability amongst medical students in a tertiary teaching hospital in South India](#)

Published On: April 08, 2017 | Pages: 025 - 028

Author(s): Haritha Sagili*, Swarup Kumar, Subitha Lakshminarayanan and Papa Dasari

Objectives: To evaluate awareness and acceptability of HPV vaccine amongst medical students in a tertiary teaching hospital. Materials and methods: This was a cross sectional descriptive study carried out in a tertiary hospital in South India in January 2015. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/jgro.000033

[Open Access](#) | [Research Article](#) | PTZAID:JGRO-3-132

[The Use of Abdominal Paracentesis in the Diagnosis of Ectopic Pregnancy in a Resource – Limited Setting](#)

Published On: March 30, 2017 | Pages: 019 - 024

Author(s): Awoleke JO* and Akintayo AA

Ectopic pregnancy remains an important cause of morbidity and mortality in the first trimester of pregnancy, and has become a major public health challenge among women in the reproductive age group in this environment. ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/jgro.000032

Review Article

[Open Access](#) | [Review Article](#) | PTZAID:JGRO-3-134

[Use of Vaginal Misoprostol Prior to Placement of an Intrauterine Device: A Review](#)

Published On: April 13, 2017 | Pages: 029 - 033

Author(s): Elizabeth Hoover, Abbey Hardy- Fairbanks and Colleen K Stockdale*

Intrauterine devices (IUD) are a highly effective and safe form of contraception. However, many individuals, particularly nulligravidas, elect against IUD use due to fear of pain associated with insertion. Misoprostol has been proposed as an agent to ease IUD insertion and decrease associated pain. ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/jgro.000034](#)

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:JGRO-3-135

[Hysterotomy for Early Placenta Percreta at 10 Weeks Gestation: A Case Report](#)

Published On: April 15, 2017 | Pages: 034 - 036

Author(s): El-Said N* and Kotb AM

Placenta percreta in early pregnancy is rare and only few cases are documented. We report a case of placenta percreta at 10 weeks gestation. Sonography revealed absence of line of demarcation between trophoblast and myometrium extending to the urinary bladder region. ...

[Abstract View](#) | [Full Article View](#) | [DOI: 10.17352/jgro.000035](#)