

In this issue

Research Article

[Open Access](#) [Research Article](#) PTZAID:JGRO-4-158

[Association of genetic polymorphisms in DNA repair genes in polycystic ovary syndrome](#)

Published On: December 27, 2018 | Pages: 044 - 048

Author(s): Sujata Dhaded* and Shailaja Dabshetty

Introduction: Polycystic ovary syndrome (PCOS) involves expression of chronic anovulation and hyperandrogenism. Role of environmental and genetic factors in PCOS is strongly supported but the genes that are positively involved in the etiology of the PCOS have not been fully investigated until now. ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/jgro.000058](https://doi.org/10.17352/jgro.000058)

[Open Access](#) [Research Article](#) PTZAID:JGRO-4-157

[The role of Umbilical Cord thickness, Interventricular Septum thickness and HbA1c levels in the prediction of Fetal Macrosomia in patients with Gestational Diabetes Mellitus](#)

Published On: December 20, 2018 | Pages: 039 - 043

Author(s): Rehab Mohamed Abdelrahman* and Mostafa Mostafa Salama

Objective: To evaluate the role of measuring umbilical cord thickness, interventricular septum thickness and HbA1c level in prediction of fetal macrosomia in patients with gestational diabetes mellitus. Methods: This prospective case-control study included 80 pregnant women. They were divided into two groups: 40 pregnant women as case group with gestational diabetes ...

[Abstract View](#) | [Full Article View](#) | DOI: [10.17352/jgro.000057](https://doi.org/10.17352/jgro.000057)

[Open Access](#) [Research Article](#) PTZAID:JGRO-4-153

[Predictors of low birth weight in antenatal women](#)

Published On: October 18, 2018 | Pages: 024 - 027

Author(s): Lakshmi S* and Rajkumar K

Background: Low birth weight is a reflection of the health care infrastructure and human development in a country. Elimination of low birth weight is essential to achieve a greater life expectancy and quality of life. There are several risk factors for low birth weight which range from maternal, nutritional, genetic and other factors. This study was carried out to evaluate ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/igro.000053

Case Report

[Open Access](#) | [Case Report](#) | PTZAID:JGRO-4-156

[Rise & Fall of Fetal Heart Rate, the Principle of Fetal Monitoring: Hypoxia Index prevents Cerebral Palsy](#)

Published On: November 29, 2018 | Pages: 036 - 038

Author(s): Kazuo Maeda* and Masaji Utsu

As FHR rises by the fetal brain excitation with fetal movement, and FHR falls by the excitation of fetal vagus nerve center with hypoxic stimulation, hypoxic grade is numerically known by the hypoxia index, which is the sum of FHR deceleration duration (min) divided by the lowest FHR (bpm), and multiplied by 100, where all of 6 cerebral palsy cases' hypoxia index were ...

[Abstract View](#) | [Full Article View](#) | DOI: 10.17352/igro.000056

[Open Access](#) | [Case Report](#) | PTZAID:JGRO-4-155

[Use of Polyglycolic Acid Mesh in Extreme Preterm Premature Rupture of Membranes \(PPROM\)](#)

Published On: November 16, 2018 | Pages: 033 - 035

Author(s): dris Koçak, Aye Özdemir and Pervin Karl*

Introduction: Extreme Preterm Premature Rupture of Membranes (PPROM) is a condition that increases maternal, fetal and neonatal morbidity and mortality to a large extent. In this study, we aimed at prolonging the gestational periods of 3 patients with extreme PPRM by way of sealing the cervical os using PGA mesh and fibrin glue to stop physical amniotic discharge. ...

[Abstract View](#)

[Full Article View](#)

[DOI: 10.17352/jgro.000055](#)

Opinion

[Open Access](#) [Opinion](#) PTZAID:JGRO-4-159

[Human zygote reconstruction by spindle, polar body or pronuclear transfer to treat repeated embryo fragmentation or embryo developmental arrest: The future is now](#)

Published On: December 31, 2018 | Pages: 049 - 051

Author(s): Konstantinos A Economou*

A common and very difficult issue to overcome as a clinical embryologist during an IVF cycle is the event of severe embryonic fragmentation or, more rarely, full developmental arrest. The current approach to bypass these phenomena is oocyte donation, but this raises several ethical concerns from the couple's side, since the offspring will bear only 50% of the biologic ...

[Abstract View](#)

[Full Article View](#)

[DOI: 10.17352/jgro.000059](#)